

*Advanced Gastrointestinal and Liver Disease*

Noah Zinkin, MD, MMSc  
775 Park Avenue, Suite 225  
Huntington, NY 11743  
(631) 923-1420

## **Clenpiq – Split Dose Prep**

\*Please remember to arrange for a responsible adult to accompany you home. If you do not have an adult escort your procedure will be cancelled and rescheduled. Taxis are not permitted.

Review the preparation schedule below for the days preceding your colonoscopy. Should you need further assistance please call 631-923-1420

### **1 Week Prior**

Go to pharmacy and fill your prescription for Prepopik. Please call our office if you did not receive a prescription.

If you take fiber supplements or vitamins containing iron, please discontinue 5 days before your appointment.

If you have diabetes, please ask us for diet and medication instructions.

If you take aspirin, plavix, Coumadin, or other blood thinners, please discuss with Dr. Zinkin whether you should continue or discontinue these prior to your procedure.

If you take Jardiance, Farxiga, Invokana or Steglatro, you must stop it 4 days before your procedure

### **3 Days Prior**

**Begin Low Fiber Diet (No fruits or vegetables. No whole wheat or high fiber. No seeds, nuts, corn or popcorn.**

**No bran or bulking agents)**

### **The Day Prior**

Begin a clear liquid diet. This includes water, clear broth or bouillon, coffee or tea, Gatorade, carbonated & non-carbonated soft drinks, Kool-Aid or other fruit-flavored drinks, strained fruit juices (no pulp), Jell-O, popsicles and hard candy. For lunch, you may have ice cream, a milkshake, or yogurt that does not contain seeds, nuts or anything solid. Drink at least 64 ounces (1/2 gallon) of clear liquid throughout the day.

## **NO RED LIQUIDS**

### **Clenpiq Instructions:**

#### **Starting at 5:30 PM the evening before your procedure:**

Drink the first bottle of Clenpiq

Drink an additional five 8 oz drinks (40 ounces total) of clear liquid over the next 3 hours.

#### **On the procedure day:**

Starting 6 hours before your procedure:

Take the second bottle of Clenpiq.

Drink an additional four 8 oz drinks (32 ounces total) of clear liquid over the next 2 hours.

Do not eat or drink anything after this.

Morning medications should be taken with a sip of water

#### **Upon departing for your procedure:**

Leave your valuables at home. Please arrange for a responsible driver to take you home. Taxis are not permitted.

Remember that no driving or work is permitted on the day of the procedure.

**Women who have had a period within 1 year will be asked for a urine specimen for a pregnancy test**